


NOAA FISHERIES West Coast Region

The West Coast Salmon Fishery



Jeromy Jording Sustainable Fishery Division, Anadromous Harvest Branch




West Coast Region

Under the Magnuson-Stevens Act, 8 regional fishery management councils develop fishery management plans specific to their regions, fisheries, and fish stocks.

Ocean Fisheries - jurisdiction

The Magnuson–Stevens Fishery Conservation and Management Act, commonly referred to as the Magnuson–Stevens Act (MSA), is the primary law governing marine U.S. coastal fisheries. Enacted in 1976, it has since been amended in line with sustainability policy.






West Coast Region

Pacific Fishery Management Council (PFMC)

The PFMC is 1 of the 8 regional management councils, and has jurisdiction over the 317,690 square mile exclusive economic zone off Washington, Oregon and California.






Annually the members of the PFMC work to develop salmon management measures to recommend to the Secretary of Commerce.


NOAA FISHERIES West Coast Region

NOAA

ARTMENT OF CO

- This is done through the National Marine Fisheries Service (NMFS).
- For PFMC, management measures are implemented by NMFS West Coast Regional offices and enforced by the NOAA Office of Law Enforcement, the 11th and 13th Coast Guard Districts, and local enforcement agencies.





PFMC Salmon season development

Pacific salmon migrate through a broad geographic range that includes rivers, streams, and coastal waters in both the United States and Canada.


Recall 1976 MSA enacted

NOAAFISHERIES

West Coast Region

 1985 Pacific salmon treaty developed

Helped ensure the respective federal governments were working with First Nations, Indian Tribes, provinces, states, and sport and commercial fishing groups to provide for shared conservation and harvest objectives.





West Coast Region

PFMC Salmon season development

Federal Court Decisions have also played key roles in defining how salmon seasons are set relative to tribal fishing rights.


- 1942 Seminole Nation v. United States
 - obligation under which the United States "has charged itself with moral obligations of the highest responsibility and trust" toward Indian tribes
- 1968 United States v. Oregon judgement
 - Case pertaining to treaty tribal fisheries in the Columbia River and stocks originating from therein
- 1974/1979/1985 United States v Washington judgement
 - Case pertaining to treaty tribal fisheries in Puget Sound and the Washington Coast and stocks originating from therein
- 1995 Parravano v. Secretary of Commerce Brown and Secretary of Interior Babbitt
 - Case pertaining to treaty tribal fisheries in the Klamath River and stocks originating from therein



West Coast Region

May October

PFMC Salmon season development



measures)

April



West Coast Region




PFMC Salmon season development

- PFMC technical committees (Salmon Technical Team [STT], Science and Statistical Committee [SSC]) collaborate to reach consensus on scientific data, methods, and approaches appropriate for recommending to the Council for management.
 - Membership mirrors the Council participating agencies
- PFMC technical committees coordinate with other committees from the other management fora
 - e.g., the *U.S. v Oregon* Technical Advisory Committee or Pacific Salmon Commission Chinook Technical Committee
- PFMC advisory subpanels containing fishery and conservation advocates provide the Council with additional input during the development process
- Public input via in-person, electronic, and paper comments are also considered



The STT uses multiple models to determine stock specific exploitation rates of different fisheries derived from recoveries of code-wire tags compared to a base period.



NOAA FISHERIES

NOAA

West Coast Region

- coveries of code-wire tags compared to a base period.

 Chinook Fishery Regulation Assessment Model (FRAM) for Chinook stocks
 - Coho FRAM for coho stocks coastwide

northwards of the mid-Oregon coast

S

- The Klamath Ocean Harvest Model (KOHM) for Klamath River fall Chinook
- The Sacramento Harvest Model (SHM) for Sacrament River fall Chinook
- The Winter Run Harvest Model (WRHM) for Sacramento River winter run Chinook



Commonalities among models:

- Have different levels of stock, fishery, area, and time strata depending on salmon fishery management need.
- Local-area total abundance not available; "catch" is an index

PFMC Salmon season development

Before the Council submits it recommendation for NMFS' adoption of annual management measures, fishery model outputs are assessed for compliance with the fishery management plan and NFMS annual guidance letter, which may specify Endangered Species Act guidance in addition to fishery management plan rules.



DOAR

West Coast Region





West Coast Region

Questions?

Thanks to:
Eric Schindler, ODFW and
Wendy Beeghley, WDFW
for providing pictures